



User Manual

MIP-3/CCL23 (MIP-3/CCL23) (Human)

Cat. No. HECCP-23013

Description:

MIP-3/CCL23 is a CC chemokine that signals through the CCR1 receptor. MIP-3 chemoattracts monocytes, resting T-lymphocytes and neutrophils, but does not chemoattract activated lymphocytes. Additionally, MIP-3 has been shown to inhibit colony formation of bone marrow myeloid immature progenitors. Alternative splicing of the MPIF1 gene results in two mRNAs that encode a short (CK β 8) and a long (CK β 81) isoform of the chemokine. CK β 8 cDNA encodes a 120 amino acid (aa) residue precursor protein with a putative 21 aa residue signal peptide that is cleaved to generate a 99 aa residue mature CK β 8 (aa 22 120). Additional N terminal processing of the 99 aa residue variant can generate a 75 aa residue CK β 8 (aa 46 120) that is significantly more active than the 99 aa residue variant.

Source:

Escherichia coli

Unit:

1 mg

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20°C. Further dilutions should be made in appropriate buffered solutions.

Formulation:

Lyophilized from a 0.2 μ m filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Molecular Weight:

Approximately 11.3 kDa, a single, non-glycosylated polypeptide chain containing 99 amino acids.

Endotoxin:

Less than 1 EU/μg of MIP-3/CCL23 as determined by LAL method.

Usage:

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

Biological Activity:

The biologically active determined by a chemotaxis bioassay using human T-lymphocytes is in a concentration of 10-50 ng/mL.

Physical Appearance:

Sterile filtered white lyophilized (freeze-dried) powder.

AA Sequence:

RVTKDAETEF MMSKLPLENP VLLDRFHATS ADCCISYTPR
SIPCSLLESYFETNSECSKP GVIFLTKKGR RFCANPSDKQ VQVCMRMLKL DTRIKTRKN

Purity:

> 97% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.