



## User Manual

**Interleukin-3 (IL-3) (Murine)**

**Cat. No. MEILP-03011**



**Description:**

IL-3 is a hematopoietic growth factor that promotes the survival, differentiation and proliferation of committed progenitor cells of the megakaryocyte, granulocyte-macrophage, erythroid, eosinophil, basophil and mast cell lineages. Produced by T cells, mast cells and eosinophils, IL-3 enhances thrombopoieses, phagocytosis, and antibody-mediated cellular cytotoxicity. Its ability to activate monocytes suggests that IL-3 may have additional immunoregulatory roles. Many of the IL-3 activities depend upon co-stimulation with other cytokines. IL-3 is species-specific, variably glycosylated cytokine.

**Source:**

*Escherichia coli*

**Unit:**

2 µg

**Reconstitution:**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20°C. Further dilutions should be made in appropriate buffered solutions.

**Formulation:**

Lyophilized from a 0.2 µm filtered solution in PBS, pH 7.4.

**Storage:**

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

**Molecular Weight:**

Approximately 14.8 kDa globular protein containing 134 amino acid residues.

**Endotoxin:**

Less than 1 EU/ $\mu$ g of IL-3 as determined by LAL method.

**Usage:**

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

**Biological Activity:**

The ED50 as determined by the dose-dependent stimulation of the proliferation of murine M-NFS-60 cells is less than 0.05 ng/mL, corresponding to a specific activity of  $> 2 \times 10^7$  units/mg.

**Physical Appearance:**

Sterile filtered white lyophilized (freeze-dried) powder.

**AA Sequence:**

DTHRLTRTLN CSSIVKEIIG KLPEPELKTD DEGPSLRNKS FRRVNLSKFV  
ESQGEVDPED RYVIKSNLQK LNCCLPTSAN DSALPGVFIR DLDDFRKKLR  
FYMVHLNDLE TVLTSRPPQP ASGSVSPNRG TVEC

**Purity:**

$> 98\%$  by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.