



User Manual

MCP-2/CCL8 (MCP-2/CCL8) (Murine)

Cat. No. MECCP-08012

Description:

MCP-2 and MCP-3 are two recently identified monocyte chemotactic proteins produced by human MG-63 osteosarcoma cells. Both MCP-2 and MCP-3 are members of the C-C family of chemokines and share 62% and 71% amino acid sequence identity, respectively, with MCP-1. MCP-3 also shares 58% amino acid identity with MCP-2. Similarly to other C-C chemokines, all three MCP proteins are monocyte chemoattractants. In addition, the three MCPs can chemoattract activated NK cells as well as CD4+ and CD8+ T lymphocytes. All three cytokines have also been shown to attract eosinophils and induce histamine secretion from basophils.

Source:

Escherichia coli

Unit:

20 µg

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20°C. Further dilutions should be made in appropriate buffered solutions.

Formulation:

Lyophilized from a 0.2 µm filtered concentrated solution in 20 mM PB, pH 7.4, 150 mM NaCl.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term

storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Molecular Weight:

Approximately 8.5 kDa, a single, non-glycosylated polypeptide chain containing 74 amino acids.

Endotoxin:

Less than 1 EU/μg of MCP-2/CCL8 as determined by LAL method.

Usage:

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

Biological Activity:

The biologically active determined by a chemotaxis bioassay using human peripheral blood monocytes is in a concentration range of 10-100 ng/mL.

Physical Appearance:

Sterile filtered white lyophilized (freeze-dried) powder.

AA Sequence:

GPDKAPVTCC FHLVCLKIPL RVLKSYERIN NIQCPMEAVV FQTKQGMSLC
VDPTQKWVSE YMEILDQKSQ ILQP

Purity:

> 97% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.